

**What is salmonellosis?**

Salmonellosis is an infection with a bacterium called *Salmonella*. Most persons infected with *Salmonella* develop diarrhea, fever and abdominal cramps 12-72 hours after infection. The illness usually lasts 4-7 days, and most persons recover without treatment; however, in some persons the diarrhea may be so severe that the patient needs to be hospitalized. In these patients, the *Salmonella* infection may spread from the intestines to the blood stream and then to other body sites and can cause death unless the person is treated promptly with antibiotics. The elderly, infants and those with impaired immune systems are more likely to have a severe illness.

**What sort of germ is *Salmonella*?**

The *Salmonella* germ is actually a group of bacteria that can cause diarrheal illness in humans. They are microscopic living creatures that pass from the feces of people or animals, to other people or other animals. There are many different kinds of *Salmonella* bacteria. *Salmonella* serotype Typhimurium and *Salmonella* serotype Enteritidis are the most common in the United States. *Salmonella* has been known to cause illness for over 100 years, having been discovered by an American scientist named Salmon, for whom they are named.

**How can *Salmonella* infections be diagnosed?**

Many different kinds of illnesses can cause diarrhea, fever or abdominal cramps. Determining that *Salmonella* is the cause of the illness depends on laboratory tests that identify *Salmonella* in the stools of an infected person. These tests are sometimes not performed unless the laboratory is instructed specifically to look for the organism. Once *Salmonella* has been identified, further testing can determine its specific type and which antibiotics could be used to treat it.

**How can *Salmonella* infections be treated?**

*Salmonella* infections usually resolve in 5 to 7 days and often do not require treatment unless the patient becomes severely dehydrated or the infection spreads from the intestines. Persons with severe diarrhea may require rehydration, often with intravenous fluids. Antibiotics are not usually necessary unless the infection spreads from the intestines, then it can be treated with ampicillin, gentamicin, trimethoprim/sulfamethoxazole (TMP/SMX) or ciprofloxacin. Unfortunately, some *Salmonella* bacteria have become resistant to antibiotics, largely as a result of the use of antibiotics to promote the growth of feed animals.

**Are there long term consequences to a *Salmonella* infection?**

Persons with diarrhea usually recover completely, although it may be several months before their bowel habits are entirely normal. A small number of persons who are infected with *Salmonella*, will go on to develop pains in their joints, irritation of the eyes and painful urination. This is called Reiter's syndrome. It can last for months or years and can lead to chronic arthritis which is difficult to treat. Antibiotic treatment does not make a difference in whether or not the person later develops arthritis.

### **How do people catch *Salmonella*?**

*Salmonella* live in the intestinal tracts of humans and other animals, including birds. *Salmonella* are usually transmitted to humans by eating foods contaminated with animal feces. Contaminated foods usually look and smell normal. Contaminated foods are often of animal origin, such as beef, poultry, milk or eggs, but all foods, including vegetables may become contaminated. Many raw foods of animal origin are frequently contaminated, but fortunately, thorough cooking kills *Salmonella*. Food may also become contaminated by the unwashed hands of an infected food handler, who forgot to wash his or her hands with soap after using the bathroom.

*Salmonella* may also be found in the feces of some pets, especially those with diarrhea, and people can become infected if they do not wash their hands after contact with these feces. Reptiles are particularly likely to harbor *Salmonella* and people should always wash their hands immediately after handling a reptile, even if the reptile is healthy. Adults should also be careful that children wash their hands after handling a reptile.

### **What can a person do to prevent this illness?**

There is no vaccine to prevent salmonellosis. Since foods of animal origin may be contaminated with *Salmonella*, people should not eat raw or undercooked eggs, poultry or meat. Raw eggs may be unrecognized in some foods such as homemade hollandaise sauce, Caesar and other salad dressings, tiramisu, homemade ice cream, homemade mayonnaise, cookie dough and frostings. Poultry and meat, including hamburgers, should be well-cooked, not pink in the middle. Persons also should not consume raw or unpasteurized milk or other dairy products. Produce should be thoroughly washed before consuming.

Cross-contamination of foods should be avoided. Uncooked meats should be kept separate from produce, cooked foods and ready-to-eat foods. Hands, cutting boards, counters, knives and other utensils should be washed thoroughly after handling uncooked foods. Hands should be washed before handling any food and between handling different food items.

People who have salmonellosis should not prepare food or pour water for others until they have been shown to no longer be carrying the *Salmonella* bacterium.

People should wash their hands after contact with animal feces. Since reptiles are particularly likely to have *Salmonella*, everyone should immediately wash their hands after handling reptiles. Reptiles (including turtles) are not appropriate pets for small children and should not be in the same house as an infant.

### **How common is salmonellosis?**

Every year, approximately 40,000 cases of salmonellosis are reported in the United States. Because many milder cases are not diagnosed or reported, the actual number of infections may be 30 or more times greater. Salmonellosis is more common in summer than winter. Children are the most likely to get salmonellosis. Young children, the elderly and the immunocompromised are the most likely to have severe infections. It is estimated that approximately 400 persons die each year with acute salmonellosis.

### **What else can be done to prevent salmonellosis?**

It is important for the public health department to know about cases of salmonellosis.

It is important for clinical laboratories to send isolates of *Salmonella* to Ohio Department of Health Laboratory (ODHL) so the specific type can be determined and compared with other *Salmonella* in the community. If many cases occur at the same time, it may mean that a restaurant, food or water supply has a problem which needs correction by the public health department.

Some prevention steps occur everyday without you thinking about it. Pasteurization of milk and treating municipal water supplies are highly effective prevention measures that have been in place for many years. In the 1970s, small pet turtles were a common source of salmonellosis in the United States, and in 1975, the sale of small turtles was halted in this country. Improvements in farm animal hygiene, in slaughter plant practices and in vegetable and fruit harvesting and packing operations may help prevent salmonellosis caused by contaminated foods. Better education of food industry workers in basic food safety and restaurant inspection procedures, may prevent cross-contamination and other food handling errors that can lead to outbreaks. Wider use of pasteurized egg in restaurants, hospitals and nursing homes is an important prevention measure. In the future, irradiation or other treatments may greatly reduce contamination of raw meat.

### **What can I do to prevent salmonellosis?**

- Cook poultry, ground beef and eggs thoroughly before eating. Do not eat or drink foods containing raw eggs or raw unpasteurized milk.
- If you are served undercooked meat, poultry or eggs in a restaurant, do not hesitate to send it back to the kitchen for further cooking.
- Wash hands, kitchen work surfaces and utensils with soap and water immediately after they have been in contact with raw meat or poultry.
- Be particularly careful with foods prepared for infants, the elderly and the immunocompromised.
- Wash hands with soap after handling reptiles and birds and after contact with pet feces.
- Avoid direct or even indirect contact between reptiles (turtles, iguanas, other lizards, snakes) and infants or immunocompromised persons.
- Do not work with raw poultry or meat and an infant (e.g. feed, change diaper) at the same time.
- Mother's milk is the safest food for young infants. Breast-feeding prevents salmonellosis and many other health problems.